

Physics 223B, Joe Kiskis

Classical groups

These are Lie groups that are subgroups of $GL(n, \mathbb{C})$.

Special groups

n vectors in an n -dimensional vector space determine a volume: $x_i = e_j x_i^j$ $i = 1, \dots, n$ gives the volume $v = \epsilon_{i_1 \dots i_n} x_1^{i_1} \dots x_n^{i_n}$. After a transformation, there are n new vectors $x'_i = e_k D_j^k x_i^j$ and a new volume $v' = \epsilon_{i_1 \dots i_n} D_{j_1}^{i_1} x_1^{j_1} \dots D_{j_n}^{i_n} x_n^{j_n}$. To get $v = v'$ for any set of vectors x_i , we need $\epsilon_{i_1 \dots i_n} D_{j_1}^{i_1} \dots D_{j_n}^{i_n} = \epsilon_{j_1 \dots j_n}$ or $\det D = 1$. Transformations satisfying this condition form a group—the special linear group $SL(n, \mathbb{C})$.

Metrics

For a more refined structure, a metric is introduced. $g(x', x) \in \mathbb{C}$ or \mathbb{R} for two vectors x and x' . g is nonsingular and linear on x $g(x', ax_1 + bx_2) = ag(x', x_1) + bg(x', x_2)$.

Bilinear metrics are also linear on x' $g(ax'_1 + bx'_2, x) = ag(x'_1, x) + bg(x'_2, x)$.

Sesquilinear metrics are still linear on x , but on x' $g(ax'_1 + bx'_2, x) = a^*g(x'_1, x) + b^*g(x'_2, x)$.

Bilinear symmetric: bilinear and $g(x', x) = g(x, x')$.

Sesquilinear symmetric: sesquilinear and $g(x', x) = g(x, x')^*$.

Bilinear antisymmetric: bilinear and $g(x', x) = -g(x, x')$.

Orthogonal groups

The field is the reals. For a bilinear symmetric metric, we can find a basis with $g_{ij} = g(e_i, e_j) = \text{diag}(1, \dots, 1, -1, \dots, -1)$ with n_+ 1's and n_- -1's. The orthogonal transformations $O(n_+, n_-, \mathbb{R})$ are those that preserve that form of the metric so that $g_{ij} = g_{kl} D_i^k D_j^l$. The further requirement that $\det D = 1$ gives the special orthogonal groups $SO(n_+, n_-, \mathbb{R})$.

Unitary groups

Now the field is the complexes. A sesquilinear symmetric can also be put in the same diagonal form. Then the transformations satisfying $g_{ij} = g_{kl} D_i^{k*} D_j^l$ form the unitary group $U(n_+, n_-)$ and with $\det D = 1$, the special unitary group $SU(n_+, n_-)$.

Symplectic groups

The field can be \mathbb{R} or \mathbb{C} . The bilinear antisymmetric metrics can be nonsingular for even n and satisfy $g_{ji} = -g_{ij}$. The symplectic groups $Sp(n, \mathbb{R} \text{ or } \mathbb{C})$ preserve a metric with this property. A common choice for g is

$$\begin{pmatrix} 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 1 & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & -1 & \cdots & 0 & 0 \\ -1 & 0 & \cdots & 0 & 0 \end{pmatrix} \quad (1)$$